

## Inclusion Criteria

1. Male or female patients, 18 years or older
2. Patients with IBD who are registered in the GEDII registry, including:
  - ❖ Patients with moderate to severe, active Crohn's disease who have not responded despite a full and adequate course of therapy with a corticosteroid and/or an immunosuppressant; or who are intolerant to or have medical contraindications for such therapies
  - ❖ Patients with fistulising active Crohn's disease who have not responded despite a full and adequate course of therapy with conventional treatment (including antibiotics, drainage and immunosuppressive therapy)
  - ❖ Patients with moderate to severe active ulcerative colitis who have had an inadequate response to conventional therapy including corticosteroids and 6-mercaptopurine (6-MP) or azathioprine (AZA), or who are intolerant to or have medical contraindications for such therapies.
3. Patients who initiated Remsima according to physician's criteria, including:
  - ❖ Anti-TNF-alfa-naïve patients
  - ❖ Patients on treatment with Remicade with stable clinical response (defined as Harvey-Bradshaw Index <5 – for CD patients; or Mayo score < 2 – for UC patients) and who switched to Remsima
4. Remsima managed according to local SMPc
5. Patients who gave their consent to be included in the GEDII Registry

## Exclusion criteria

1. Patients who are not eligible for anti-TNF-alfa therapy
2. Patients who are being treated with any investigational agent
3. Patients who are not willing to comply with routine clinical appointments

## Laboratory Testing

### Serum samples:

- For all time points, marked in the chronogram, collect 5 mL of blood to a serum gel tube
- Blood must be left for 1 hour at room temperature to assure coagulation
- Centrifuge at 3000 rpm, for 10 min
- Collect serum to two cryovials
- Identify cryovials with the label provided in the kit
- Freeze and store serum samples at -20°C until refrigerated transport is provided

### Fecal samples:

- Each sample should be collected with the EASYSample collection Kit
- The tube must be identified with the label provided in the kit
- Send samples to the central laboratory within 48 hours
- **Contact GEDII for transport**

## Study Duration

The overall duration of the study is three years (1 year of recruitment + 2-year observation period).

## Study Timelines

The study is expected to start during 2015. Study closure is expected to occur 2nd Quarter of 2018.

**For any study related questions, please contact:**

**Sandra Dias**

**E-mail:** [gedi@med.up.pt](mailto:gedi@med.up.pt)

**Phone number:** 912583273

## REMREGISTER

**Title:** A study in the real-world practice to evaluate the impact of biosimilar infliximab (Remsima) in clinical outcomes in patients with inflammatory bowel diseases: a 2-year longitudinal analysis from the GEDII Registry



**Study sponsor:** Grupo de Estudo da Doença Inflamatória Intestinal (GEDII)

**Name of Scientific Coordinator:** Prof. Fernando Magro

## Primary Objectives

- To evaluate the impact of Remsima in inducing clinical response among biologic-naïve patients with IBDs registered in the GEDII Registry followed for two years.
- To evaluate the impact of Remsima in inducing clinical remission among biologic-naïve patients with IBDs registered in the GEDII Registry followed for two years.
- To evaluate the impact of Remsima in promoting mucosal healing, among biologic-naïve patients registered in the GEDII Registry followed for two years.
- To evaluate the impact of Remsima in promoting biomarkers remission by normalization of calprotectin.

**CHRONOGRAM A (applicable for CD or fistulising CD patients)**

|  | Infliximab        |     |  |      |      |      |      |      |      |      |      |      |      |      |       |                                   |
|--|-------------------|-----|--|------|------|------|------|------|------|------|------|------|------|------|-------|-----------------------------------|
|  | Induction period* |     | Maintenance period – data collection time point (24 month follow up) |      |      |      |      |      |      |      |      |      |      |      |       |                                   |
| Information to be collected  | Day1 or basal     | W 2 | W 6  | W 14 | W 22 | W 30 | W 38 | W 46 | W 54 | W 62 | W 70 | W 78 | W 86 | W 94 | W 102 | Or completion of Fup <sup>3</sup> |
| Date of Birth  | X <sup>4</sup>    |     |  |      |      |      |      |      |      |      |      |      |      |      |       |                                   |
| Sex  | X <sup>4</sup>    |     |  |      |      |      |      |      |      |      |      |      |      |      |       |                                   |
| Height   | X <sup>4</sup>    |     |  |      |      |      |      |      |      |      |      |      |      |      |       |                                   |
| Weight   | X                 | X   | X  | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X     | X                                 |
| BMI  | X                 | X   | X  | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X     | X                                 |
| Smoking status   | X <sup>4</sup>    |     |  |      |      |      |      |      |      |      |      |      |      |      |       |                                   |
| Medical history  | X <sup>4</sup>    |     |  |      |      |      |      |      |      |      |      |      |      |      |       |                                   |
| Comorbidities  | X                 | X   | X  | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X     | X                                 |
| Disease presentation   | X <sup>4</sup>    |     |  |      |      |      |      |      |      |      |      |      |      |      |       |                                   |
| Diagnosis CD, location, steroid, behaviour, prognostic classification. | X <sup>4</sup>    |     |  |      |      |      |      |      |      |      |      |      |      |      |       |                                   |
| Clinical activity (HBI)  | X                 | X   | X  | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X     | X                                 |
| Nr of draining fistulae (fistulising CD)                               | X                 | X   | X  | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X     | X                                 |
| PDAI (perianal CD)   | X                 | X   | X  | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X     | X                                 |
| Dose of Remsima administered   | X                 | X   | X  | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X     | X                                 |
| Endoscopy  | X                 |     |  |      |      |      |      |      | X    |      |      |      |      |      | X     | X                                 |
| Concomitant therapies  | X                 | X   | X  | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X     | X                                 |
| Routine laboratory parameters  | X                 | X   | X  | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X     | X                                 |
| Serum infliximab levels <sup>1</sup>                                   | X                 | X   | X  | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X     | X                                 |
| Anti-drug antibodies <sup>1</sup>                                      | X                 | X   | X  | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X     | X                                 |
| Fecal sample (calprotectin levels) <sup>2</sup>                        | X                 | X   | X  | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X     | X                                 |
| Adverse reactions  | X                 | X   | X  | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X     | X                                 |
| Use of health resources  | X                 | X   | X  | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X     | X                                 |

\*Data related with induction period **will only be collected** for biologic-naïve patients stating Remsima. Data related with maintenance period is applicable both for biologic-naïve patients and patients who switched from Remicade. W = week, HBI = Harvey Bradshaw Index, PDAI, Perianal Disease Activity Index.  
<sup>1</sup>Blood sample to be collected prior to each infusion of Remsima at the scheduled appointment. Sample will be analysed by Central Laboratory.  
<sup>2</sup>Stool sample to be collected and sent for Centro Laboratory  
<sup>3</sup>Patients in maintenance with Remsima at study inclusion will be followed every 8 weeks (or every 6 weeks, if required) until completing the 24-month follow up period;  
<sup>4</sup>Basal data to be collected, regardless of the patient's treatment status at study inclusion

**CHRONOGRAM B (applicable for UC patients)**

|  | Infliximab        |     |  |      |      |      |      |      |      |      |      |      |      |      |       |                                   |
|--|-------------------|-----|--|------|------|------|------|------|------|------|------|------|------|------|-------|-----------------------------------|
|  | Induction period* |     | Maintenance period – data collection time point (24 month follow up) |      |      |      |      |      |      |      |      |      |      |      |       |                                   |
| Information to be collected  | Day1 or basal     | W 2 | W 6  | W 14 | W 22 | W 30 | W 38 | W 46 | W 54 | W 62 | W 70 | W 78 | W 86 | W 94 | W 102 | Or completion of Fup <sup>3</sup> |
| Date of Birth  | X <sup>4</sup>    |     |  |      |      |      |      |      |      |      |      |      |      |      |       |                                   |
| Sex  | X <sup>4</sup>    |     |  |      |      |      |      |      |      |      |      |      |      |      |       |                                   |
| Height   | X <sup>4</sup>    |     |  |      |      |      |      |      |      |      |      |      |      |      |       |                                   |
| Weight   | X                 | X   | X  | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X     | X                                 |
| BMI  | X                 | X   | X  | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X     | X                                 |
| Smoking status   | X <sup>4</sup>    |     |  |      |      |      |      |      |      |      |      |      |      |      |       |                                   |
| Medical history  | X <sup>4</sup>    |     |  |      |      |      |      |      |      |      |      |      |      |      |       |                                   |
| Comorbidities  | X                 | X   | X  | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X     | X                                 |
| Disease presentation   | X <sup>4</sup>    |     |  |      |      |      |      |      |      |      |      |      |      |      |       |                                   |
| Diagnosis CD, location, steroid, behaviour, prognostic classification. | X <sup>4</sup>    |     |  |      |      |      |      |      |      |      |      |      |      |      |       |                                   |
| Clinical activity (Partial Mayo)                                       | X                 | X   | X  | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X     | X                                 |
| Dose of Remsima administered   | X                 | X   | X  | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X     | X                                 |
| Endoscopy  | X                 |     |  | X    |      |      |      |      | X    |      |      |      |      |      | X     | X                                 |
| Concomitant therapies  | X                 | X   | X  | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X     | X                                 |
| Routine laboratory parameters  | X                 | X   | X  | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X     | X                                 |
| Serum infliximab levels <sup>1</sup>                                   | X                 | X   |  | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X     | X                                 |
| Anti-drug antibodies <sup>1</sup>                                      | X                 | X   |  | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X     | X                                 |
| Fecal sample (calprotectin levels) <sup>2</sup>                        | X                 | X   | X  | X    | X    |      |      | X    |      |      | X    |      | X    | X    | X     | X                                 |
| Adverse reactions  | X                 | X   | X  | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X     | X                                 |
| Use of health resources  | X                 | X   | X  | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X    | X     | X                                 |

\* Data related with induction period **will only be collected** for biologic-naïve patients starting Remsima. Data related with maintenance period is applicable both to biologic-naïve patients and patients who switched from Remicade.  
<sup>1</sup> Blood sample to be collected prior to each infusion of Remsima at the scheduled appointment. Sample will be analyzed by Central Laboratory.  
<sup>2</sup> Stool sample to be collected and sent for Central Laboratory  
<sup>3</sup> Patients in maintenance with Remsima at study inclusion will be followed every 8 weeks (or every 6 weeks, if required) until completing the 24-month follow up period;  
<sup>4</sup> Basal data to be collected, regardless of the patient's treatment status at study inclusion. W = week